


# Introduction to Modern Virology

FOURTH EDITION

BASIC  
MICROBIOLOGY  
SERIES

N J Dimmock  
S B Primrose



*b*

Blackwell  
Science

Best Available Copy

© 1974, 1980, 1987, 1994 by  
Blackwell Science Ltd  
Editorial Offices:  
Osney Mead, Oxford OX2 0EL  
25 John Street, London WC1N 2BL  
23 Ainslie Place, Edinburgh EH3 6AJ  
236 Main Street, Cambridge  
Massachusetts 02142, USA  
54 University Street, Carlton  
Victoria 3053, Australia

Other Editorial Offices:  
Amette Blackwell SA  
1, rue de Lille, 75007 Paris  
France

Blackwell Wissenschafts-Verlag GmbH  
Kurfürstendamm 57  
10707 Berlin, Germany

Blackwell MZV  
Feldgasse 13, A-1238 Wien  
Austria

All rights reserved. No part of this  
publication may be reproduced, stored in a  
retrieval system, or transmitted, in any  
form or by any means, electronic,  
mechanical, photocopying, recording or  
otherwise, except as permitted by the UK  
Copyright, Designs and Patents Act 1988,  
without the prior permission of the  
copyright owner.

First published 1974  
Reprinted 1978  
Second edition 1980  
Third edition 1987  
Fourth edition 1994

Set by Setrite Typesetters, Hong Kong  
Printed and bound in Great Britain  
at the University Press, Cambridge

#### DISTRIBUTORS

Marston Book Services Ltd  
PO Box 87  
Oxford OX2 0DT  
(Orders: Tel: 01865 791155  
Fax: 01865 791927  
Telex: 837515)

#### USA

Blackwell Science, Inc.  
238 Main Street  
Cambridge, MA 02142  
(Orders: Tel: 800 759-6102  
617 876-7000)

#### Canada

Oxford University Press  
70 Wynford Drive  
Don Mills  
Ontario M3C 1J9  
(Orders: Tel: 416 441-2941)

#### Australia

Blackwell Science Pty Ltd  
54 University Street  
Carlton, Victoria 3053  
(Orders: Tel: 03 347-5552)

A catalogue record for this title  
is available from the British Library  
ISBN 0-632-03403-3

Library of Congress  
Cataloging-in-Publication Data

Dimmock, N. J.  
Introduction to modern virology/  
N. J. Dimmock, S. B. Primrose. — 4th ed.  
p. cm.

Includes bibliographical references  
and index.

ISBN 0-632-03403-3

1. Virology. I. Primrose, S. B.

II. Title. III. Series.

[DNLM: 1. Virus Diseases.

2. Viruses. QW 160 D582i 1994]

QR360.D56 1994

616'.0194—dc20

Best Available Copy

## 184 Chapter 11

**The assembly of enveloped viruses**

A large number of viruses, particularly viruses infecting animals, have a lipid envelope as an integral part of their structure. These include many animal viruses (herpes-, toga-, oncorna-, orthomyxo-, paramyxo-, corona-, arena-, pox- and irido-), the tomato spotted wilt virus group and rhabdoviruses of plants and three families of bacteriophage (cortico-, plasma- and cystoviruses) [see Table 3.2 and Chapter 21]. Herpesviruses replicate in the cell nucleus, although the viral proteins are synthesized in the cytoplasm and transported back into the nucleus. After assembly of the nucleocapsid, the virus buds off from the nuclear membrane and thus becomes enveloped. Prior to the budding process, the membrane is modified by incorporation of viral-specified proteins, which are subsequently glycosylated. Very little is known about the assembly of the lipid-containing phages and the iridoviruses, but it appears that the envelope is not incorporated by a budding process. In this respect, they resemble poxviruses, whose morphogenesis has been studied extensively by electron microscopy of infected cells. In thin sections, particles initially appear as crescent-shaped objects within specific areas of cytoplasm, called 'factories', and even at this stage they appear to contain the trilaminar membrane which forms the envelope. The crescents are then completed into spherical structures. DNA is added, and then the external surface undergoes a number of modifications to yield mature virions.

The majority of enveloped viruses acquire their envelope by budding from the plasma membrane or one of the internal cytoplasmic membranes (Fig. 11.13). Four events leading to the maturation have been identified. First, the nucleocapsids form in the cytoplasm. Secondly, patches of cellular membrane incorporate viral glycoproteins, which are transmembrane proteins. Thirdly, these are focused by interaction of their cytoplasmic tails with the nucleocapsid (or intermediary matrix protein), which becomes aligned along the inner surface of the modified membrane, and finally the virion is formed by budding. It is remarkable that during the budding process host membrane proteins are excluded from viral particles (retroviruses are the exception), although the bulk of the lipid in the envelope is derived from the host's normal complement of lipid.

Viruses budding into the endoplasmic reticulum are released to the exterior via the Golgi complex. First, the virus particle buds into a vesicle, which moves to and fuses with Golgi complex. Then, following the normal route of transport of cellular proteins, the virus moves through the Golgi and buds from the concave surface into another vesicle, which is transported to the plasma membrane. Fusion of the vesicle with the plasma membrane releases the particle to the exterior of the cell. Viruses which bud from the plasma membrane are automatically released when the budding process is complete.

## The process of infection: IV 185




Fig. 11.13 Sites of maturation of various enveloped viruses. F, fusion of a vesicle with a membrane.

One complication is that many differentiated cells *in vivo* are polarized, meaning that they carry out different functions with their outer (apical) surface and their inner (basal) surface. For instance, cells lining kidney tubules are responsible for regulating  $\text{Na}^+$  ion concentration. They transport  $\text{Na}^+$  ions via their apical surface from blood to the cytoplasm and then expel them from the basal surface into urine. Some cell lines, such as Madin-Darby canine kidney (MDCK) cells, retain this property in culture, but this can only be demonstrated when they form confluent monolayers and tight junctions between cells. The latter serve to separate and define properties of the apical and basal surfaces. These properties reside in cellular proteins which have migrated directionally to one or other surface. This happens too with some viral proteins. The lipid composition of a polarized cell is also distributed asymmetrically between apical and basal surfaces. For instance, if a cell is dually infected with a rhabdovirus and an orthomyxovirus,